

Least Squares Regression Line

Q1, (OCR 4767, Jan 2007, Q1i-iii)

<p>(i)</p>	<p><math>\bar{t} = 112.8, \bar{v} = 0.6</math></p> $b = \frac{S_{vt}}{S_{vv}} = \frac{405.2 - 3 \times 564 / 5}{2.20 - 3^2 / 5} = \frac{66.8}{0.4} = 167$ <p>OR <math>b = \frac{405.2 / 5 - 0.6 \times 112.8}{2.20 / 5 - 0.6^2} = \frac{13.36}{0.08} = 167</math></p> <p>hence least squares regression line is:</p> $t - \bar{t} = b(v - \bar{v})$ $\Rightarrow t - 112.8 = 167(v - 0.6)$ $\Rightarrow t = 167v + 12.6$	<p>B1 for <math>\bar{t}</math> and <math>\bar{v}</math> used (SOI)</p> <p>M1 for attempt at gradient (b)</p> <p>A1 for 167 CAO</p> <p>M1 for equation of line</p> <p>A1 FT</p>	<p><b>5</b></p>
<p>(ii)</p>	<p>(A) For 0.5 litres, predicted time = = <math>167 \times 0.5 + 12.6 = 96.1</math> seconds</p> <p>(B) For 1.5 litres, predicted time = = <math>167 \times 1.5 + 12.6 = 263.1</math> seconds</p> <p>Any valid relevant comment relating to each prediction such as eg: 'First prediction is fairly reliable as it is interpolation and the data is a good fit' 'Second prediction is less certain as it is an extrapolation'</p>	<p>M1 for at least one prediction attempted</p> <p>A1 for both answers (FT their equation if <math>b &gt; 0</math>) NB for reading predictions off the graph only award A1 if accurate to nearest whole number</p> <p>E1 (first prediction) E1 (second prediction)</p>	<p><b>4</b></p>
<p>(iii)</p>	<p>The <math>v</math>-coefficient is the number of additional seconds required for each extra litre of water</p>	<p>E1 for indication of rate wrt <math>v</math> E1 <i>dep</i> for specifying its units</p>	<p><b>2</b></p>

**Q2, (Jan 2008, Q1i-iii)**

<b>(i)</b>	<p><math>x</math> is independent, <math>y</math> is dependent since the values of <math>x</math> are chosen by the student but the values of <math>y</math> are dependent on <math>x</math></p>	<p>B1 E1 dep E1 dep</p>	<b>3</b>
<b>(ii)</b>	<p><math>\bar{x} = 2.5, \bar{y} = 80.63</math></p> $b = \frac{S_{xy}}{S_{xx}} = \frac{2530.3 - 30 \times 967.6/12}{90 - 30^2/12} = \frac{111.3}{15} = 7.42$ <p>OR <math>b = \frac{2530.3/12 - 2.50 \times 80.63}{90/12 - 2.50^2} = \frac{9.275}{1.25} = 7.42</math></p> <p>Hence least squares regression line is:  <math>y - \bar{y} = b(x - \bar{x})</math>  <math>\Rightarrow y - 80.63 = 7.42(x - 2.5)</math>  <math>\Rightarrow y = 7.42x + 62.08</math></p>	<p>B1 for <math>\bar{x}</math> and <math>\bar{y}</math> used (SOI)</p> <p>M1 for attempt at gradient (<math>b</math>) A1 for 7.42 <b>cao</b></p> <p>M1 for equation of line</p> <p>A1 FT (<math>b &gt; 0</math>) for complete equation</p>	<b>5</b>
<b>(iii)</b>	<p>(A) For <math>x = 1.2</math>, predicted growth  <math>= 7.42 \times 1.2 + 62.08 = 71.0</math></p> <p>(B) For <math>x = 4.3</math>, predicted growth  <math>= 7.42 \times 4.3 + 62.08 = 94.0</math></p> <p>Valid relevant comments relating to the predictions such as :          Comment re interpolation/extrapolation          Comment relating to the fact that <math>x = 4.3</math> is only just beyond the existing data.          Comment relating to size of residuals near each predicted value (need not use word 'residual')</p>	<p>M1 for at least one prediction attempted. A1 for both answers (FT their equation if <math>b &gt; 0</math>)</p> <p>E1 (first comment)</p> <p>E1 (second comment)</p>	<b>4</b>


**Q3, (OCR 4732, Jan 2005, Q9)**

<b>(i)</b>	<p><math>x</math> is independent, <math>y</math> is dependent since the values of <math>x</math> are chosen by the student but the values of <math>y</math> are dependent on <math>x</math></p>	<p>B1 E1 dep E1 dep</p>	<b>3</b>
<b>(ii)</b>	<p><math>\bar{x} = 2.5, \bar{y} = 80.63</math> <math>b = \frac{S_{xy}}{S_{xx}} = \frac{2530.3 - 30 \times 967.6/12}{90 - 30^2/12} = \frac{111.3}{15} = 7.42</math> OR <math>b = \frac{2530.3/12 - 2.50 \times 80.63}{90/12 - 2.50^2} = \frac{9.275}{1.25} = 7.42</math> Hence least squares regression line is: <math>y - \bar{y} = b(x - \bar{x})</math> <math>\Rightarrow y - 80.63 = 7.42(x - 2.5)</math> <math>\Rightarrow y = 7.42x + 62.08</math></p>	<p>B1 for <math>\bar{x}</math> and <math>\bar{y}</math> used (SOI) M1 for attempt at gradient (<math>b</math>) A1 for 7.42 <b>cao</b> M1 for equation of line A1 FT (<math>b &gt; 0</math>) for complete equation</p>	<b>5</b>
<b>(iii)</b>	<p>(A) For <math>x = 1.2</math>, predicted growth <math>= 7.42 \times 1.2 + 62.08 = 71.0</math> (B) For <math>x = 4.3</math>, predicted growth <math>= 7.42 \times 4.3 + 62.08 = 94.0</math>  Valid relevant comments relating to the predictions such as : Comment re interpolation/extrapolation Comment relating to the fact that <math>x = 4.3</math> is only just beyond the existing data. Comment relating to size of residuals near each predicted value (need not use word 'residual')</p>	<p>M1 for at least one prediction attempted. A1 for both answers (FT their equation if <math>b &gt; 0</math>)  E1 (first comment) E1 (second comment)</p>	<b>4</b>


**Q4, (OCR 4732, Jan 2009, Q2)**

<b>(i)(a)</b>	$\frac{8736.9 - \frac{202 \times 245.3}{7}}{7300 - \frac{202^2}{7}} \text{ or } \frac{1658.24}{1470.86}$ <p><math>= 1.127\dots</math> (= 1.13 <b>AG</b>)</p>	<p>M1  A1 2</p>	<p>correct sub in any correct formula for <math>b</math> eg <math>\frac{236.8921}{210.1249}</math>  must see 1.127... ; 1.127.. alone: M1A1</p>
<b>(b)</b>	<p><math>y - \frac{245.3}{7} = 1.13(x - \frac{202}{7})</math> <math>y = 1.1x + 2.5</math> (or 2.4) or <math>y = 1.13x + 2.43</math></p>	<p>M1 A1 2</p>	<p>or <math>a = \frac{245.3}{7} - 1.13 \times \frac{202}{7}</math> 2 sfs suff. (exact: <math>y = 1.127399\dots x + 2.50934\dots</math>)</p>
<b>(ii)(a)</b>	<p><math>(1.1(\dots) \times 30 + 2.5(\dots)) = 35.5</math> to <math>36.5</math></p>	<p>B1f 1</p>	
<b>(b)</b>	<p><math>(1.1(\dots) \times 100 + 2.5(\dots)) = 112.4</math> to <math>115.6</math></p>	<p>B1f 1</p>	
<b>(iii)</b>	<p>(a) Reliable  (b) Unreliable because extrapolated</p>	<p>B1  B1 2</p>	<p>Both reliable: B1  Ignore extras  (a) more reliable than (b) B1 because (a) within data or (b) outside data B1</p>

**Q5, (OCR 4767, Jan 2013, Q1)**

<p>(i)</p>		<p>G1</p> <p>G2,1,0</p> <p>[3]</p>	<p>G1 For axes suitably labelled with some indication of <b>linear</b> scale provided.</p> <p>G2 for points plotted correctly. G1 if 4 points plotted correctly. G0 if two or more incorrectly plotted/omitted points.</p> <p>Special Case SC1 for points visibly correct on axes where no indication of scale has been provided.</p>	<p>Allow <math>x</math> &amp; <math>y</math> Allow axes reversed.</p>
<p>(ii)</p>	<p>Thickness is the independent variable since the values of ‘Thickness’ are not subject to random variation, but are determined by the manufacturer.</p>	<p>E1</p> <p>[1]</p>	<p>Allow explanations referring to thickness being “controlled” by the manufacturer. Allow equivalent interpretations.</p>	
<p>(iii)</p>	<p><math>\bar{t} = 60, \bar{h} = 1.548</math></p> $b = \frac{S_{th}}{S_{tt}} = \frac{546.8 - (300 \times 7.74 / 5)}{22000 - 300^2 / 5} = \frac{82.4}{4000} = 0.0206$	<p>B1</p> <p>M1*</p> <p>A1</p>	<p>For <math>\bar{t}</math> and <math>\bar{h}</math> used. SOI (e.g. can be implied by <math>b = 0.0206</math>)</p> <p>For attempt at <b>calculating</b> gradient (<math>b</math>) for <math>h</math> on <math>t</math>.</p> <p>For 0.0206 cao</p>	
	<p>OR <math>b = \frac{546.8 / 5 - (60 \times 1.548)}{22000 / 5 - 60^2} = \frac{16.48}{800} = 0.0206</math></p> <p>hence least squares regression line is:</p> $h - \bar{h} = b(t - \bar{t})$			

**Q6, (OCR 4767, Jun 2015, Q1i,ii,iv,v)**

<p>(i)</p>		<p>G1* Both axes labeled (allow <math>t</math> and <math>y</math>) with indication of scale</p> <p>G1<sub>dep</sub>* for values of time BOD if (0,0) not clearly visible</p> <p>G1<sub>dep</sub>* for values of average growth BOD if (0,0) not clearly visible.</p> <p>BOD if confusion arises from points plotted for part (v).</p>	<p>Allow axes interchanged Condone <math>x</math> for <math>t</math></p> <p>(evenly spaced)</p> <p>visually correct</p> <p>SC1 for points having the correct distribution and G0* awarded.</p> <p>Line through origin should appear but this is rewarded in part (v)</p>
<p>(ii)</p>	<p><math>\bar{t} = 1.5, \bar{y} = 32</math></p> $b = \frac{S_{yt}}{S_{tt}} = \frac{490 - (224 \times 10.5 / 7)}{22.75 - 10.5^2 / 7} = \frac{154}{7} = 22$ <p>OR <math>b = \frac{490 / 7 - (32 \times 1.5)}{22.75 / 7 - 1.5^2} = \frac{22}{1} = 22</math></p> <p>hence least squares regression line is:</p> $y - \bar{y} = b(t - \bar{t})$ $\Rightarrow y - 32 = 22(t - 1.5)$ $\Rightarrow y = 22t - 1$	<p>B1 For <math>\bar{t}</math> and <math>\bar{y}</math> seen or implied by final answer.</p> <p>M1 For attempt at gradient (<math>b</math>)</p> <p>A1 For 22 cao</p> <p>M1 For equation of line</p> <p>A1 CAO</p>	<p>Seen either in calculating <math>b</math> or in forming the equation of the line.</p> <p>Correct structure needed. See additional notes. FT their <math>\bar{t}</math> and <math>\bar{y}</math> for M1</p> <p>With their <math>b &gt; 0, \bar{t}</math> and <math>\bar{y}</math></p> <p>A0 for <math>y = 22x - 1</math></p>

[3]

[5]

(iv)		$(22 \times 5) - 1 = 109$  Likely to be <b>unreliable as extrapolation</b> (oe)	B1 B1 [2]	Estimate calculated using equation	FT their equation
(v)		$a = \frac{490}{22.75} = 21.538\dots = 21.5$ (3 s.f.) Equation is $y = 21.5t$ Line plotted on diagram	M1 A1 A1 A1  [4]	Allow $y = 21.54t$ CAO For line correctly plotted CAO A0 if axes not scaled or $a \neq 21.5$ to 3 sf	Allow $y = (280/13)t$ Through (0,0) and between (3, 64) and (3,65)

**Q7, (OCR 4732, Jun 2016, Q2ii)**

ii	a	'Increased' <u>and</u>  Positive gradient or positive coeff of $n$ or 'Output goes up by 0.6 each month'  Both needed	B1 [1]	'Increased' <u>and</u>  values of $z$ shown as follows: at least 6 values or 1st and last values or 1st, or 2nd or 3rd or 4th <u>and</u> 9th or 10th or 11th or 12th ie 17.6 or 18.2 or 18.8 or 19.4 <u>and</u> 22.4 or 23 or 23.6 or 24.2	'Increased' <u>and</u>  'Value of $0.6n$ increases as $n$ increases'	<u>NOT:</u> 'Increased' and 'Value of $z$ incr as $n$ incr' ' $z$ incr as no. of mths incr'
ii	b	$\bar{n} = 6.5$ or $\frac{78}{12}$ oe seen $\bar{z} = 0.6 \times 6.5 + 17$ alone, eg not $+12$ or $17 = \bar{z} - 0.6 \times 6.5$ oe  $\bar{z} = 20.9$	B1 M1 A1 [3]	or $(0.6 \times 1 + 17 + 0.6 \times 2 + 17 + \dots + 0.6 \times 12 + 17) \div 12$ oe or '250.8' $\div 12$ M1 ft their '6.5' only if comes from $\dots + 12$ cao	Long method, all correct terms seen and $+12$ M1  NB ans 20.9 may not score the B1	
ii	c	Total output = "20.9" $\times 12$  251 (3 sf)	M1 A1f [2]	or $0.6 \times 1 + 17 + 0.6 \times 2 + 17 + \dots + 0.6 \times 12 + 17$ oe or eg $\frac{88}{5} + \frac{91}{5} + \dots + \frac{121}{5}$ oe ft their (ii)(b)	Long method, all correct terms seen  Not ISW, eg 25100 scores A0, even if 251 seen	
			<b>10</b>			