

Exponential and Logarithmic Equations Exam Questions MS

Q1, (OCR 4722, Jun 2009, Q3)

$$\log 7^x = \log 2^{x+1}$$

$$x \log 7 = (x+1) \log 2$$

$$x(\log 7 - \log 2) = \log 2$$

$$x = 0.553$$

- M1 Introduce logarithms throughout, or equiv with base 7 or 2
- M1 Drop power on at least one side
- A1 Obtain correct linear equation (allow with no brackets)
- M1 **Either** expand bracket and attempt to gather x terms,
or deal correctly with algebraic fraction
- A1 **5** Obtain $x = 0.55$, or rounding to this, with no errors seen

5

Q2, (OCR 4722, Jun 2010, Q8)

a $\log 5^{3w-1} = \log 4^{250}$

$$(3w-1) \log 5 = 250 \log 4$$

$$3w-1 = \frac{250 \log 4}{\log 5}$$

$$w = 72.1$$

- M1* Introduce logarithms throughout
- M1* Use $\log a^b = b \log a$ at least once
- A1 Obtain $(3w-1) \log 5 = 250 \log 4$ or equiv
- M1d* Attempt solution of linear equation
- A1 **5** Obtain 72.1, or better

b $\log_x \frac{5y+1}{3} = 4$

$$\frac{5y+1}{3} = x^4$$


$$5y+1 = 3x^4$$

$$y = \frac{3x^4 - 1}{5}$$

- M1 Use $\log a - \log b = \log \frac{a}{b}$ or equiv
- M1 Use $f(y) = x^4$ as inverse of $\log_x f(y) = 4$
- M1 Attempt to make y the subject of $f(y) = x^4$
- A1 **4** Obtain $y = \frac{3x^4 - 1}{5}$, or equiv

9

Q3, (OCR 4722, Jun 2008, Q8)



- M1** Attempt sketch of exponential graph (1st quad)
- if seen in 2nd quad must be approx correct
- A1** Correct graph in both quadrants
- B1** State or imply (0, 2) only

3

(ii) $8^x = 2 \times 3^x$
 $\log_2 8^x = \log_2 (2 \times 3^x)$

$x \log_2 8 = \log_2 2 + x \log_2 3$

$3x = 1 + x \log_2 3$

$x(3 - \log_2 3) = 1$, hence $x = \frac{1}{3 - \log_2 3}$ **A.G.**

OR $8^x = 2 \times 3^x$

$2^{3x} = 2 \times 3^x$

$2^{(3x-1)} = 3^x$

$\log_2 2^{(3x-1)} = \log_2 3^x$

$(3x - 1) \log_2 2 = x \log_2 3$


$x(3 - \log_2 3) = 1$, hence $x = \frac{1}{3 - \log_2 3}$ **A.G.**

- M1** Form equation in x and take logs (to any consistent base, or no base) – could use \log_8
- M1** Use $\log a^b = b \log a$
- M1** Use $\log ab = \log a + \log b$, or equiv with $\log^{a/b}$
- M1** Use $\log_2 8 = 3$
- A1** Show given answer correctly

- M1** Use $8^x = 2^{3x}$
- M1** Attempt to rearrange equation to $2^k = 3^x$
- M1** Take logs (to any base)
- M1** Use $\log a^b = b \log a$
- A1** Show given answer correctly

5

Q4, (OCR 4722, Jan 2010, Q9)



- M1** Reasonable graph in both quadrants
- A1** Correct graph in both quadrants

B1 3 State or imply (0, 6)

(ii) $9^x = 150$

$x \log 9 = \log 150$

$x = 2.28$

- M1** Introduce logarithms throughout, or equiv with \log_9
- M1** Use $\log a^b = b \log a$ and attempt correct method to find x
- A1** 3 Obtain $x = 2.28$

(iii) $6 \times 5^x = 9^x$

$\log_3 (6 \times 5^x) = \log_3 9^x$

$\log_3 6 + x \log_3 5 = x \log_3 9$

$\log_3 3 + \log_3 2 + x \log_3 5 = 2x$

$x(2 - \log_3 5) = 1 + \log_3 2$

$x = \frac{1 + \log_3 2}{2 - \log_3 5}$ **A.G.**

- M1** Form eqn in x and take logs throughout (any base)
- M1** Use $\log a^b = b \log a$ correctly on $\log 5^x$ or $\log 9^x$ or legitimate combination of these two
- M1** Use $\log ab = \log a + \log b$ correctly on $\log (6 \times 5^x)$ or $\log 6$
- M1** Use $\log_3 9 = 2$ or equiv (need base 3 throughout that line)
- A1** 5 Obtain $x = \frac{1 + \log_3 2}{2 - \log_3 5}$ convincingly
(inc base 3 throughout)

11

Q5, (OCR 4722 Jan 2009, Q8)

<p>(a)(i) $\log_a xy = p + q$</p>	<p>B1 1 State $p + q$ cwo</p>
<hr style="border-top: 1px dashed black;"/>	
<p>(ii) $\log_a \left(\frac{a^2 x^3}{y}\right) = 2 + 3p - q$</p>	<p>M1 Use $\log a^b = b \log a$ correctly at least once</p> <p>M1 Use $\log \frac{a}{b} = \log a - \log b$ correctly</p> <p>A1 3 Obtain $2 + 3p - q$</p>
<hr style="border-top: 1px dashed black;"/>	
<p>(b)(i) $\log_{10} \frac{x^2 - 10}{x}$</p>	<p>B1 1 State $\log_{10} \frac{x^2 - 10}{x}$ (with or without base 10)</p>
<hr style="border-top: 1px dashed black;"/>	
<p>(ii) $\log_{10} \frac{x^2 - 10}{x} = \log_{10} 9$</p> <p>$\frac{x^2 - 10}{x} = 9$</p> <p>$x^2 - 9x - 10 = 0$</p> <p>$(x - 10)(x + 1) = 0$</p> <p>$x = 10$</p>	<p>B1 State or imply that $2 \log_{10} 3 = \log_{10} 3^2$</p> <p>M1 Attempt correct method to remove logs</p> <p>A1 Obtain correct $x^2 - 9x - 10 = 0$ aef, no fractions</p> <p>M1 Attempt to solve three term quadratic</p> <p>A1 5 Obtain $x = 10$ only</p>

Q6, (OCR 4722, Jan 2013, Q8)

(i)		Translation of 3 units in positive x -direction	B1	State translation
			B1	State or imply 3 units in positive x -direction
(ii)		$a = 8$	B1 [1]	State 8
(iii)		$b - 3 = 2^{1.8}$ $b = 6.48$	B1	State or imply $b - 3 = 2^{1.8}$
			B1 [2]	Obtain 6.48, or better
(iv)		$\log_2 c - \log_2(c - 3) = 4$ $\log_2 {}^c/c-3 = 4$ ${}^c/c-3 = 2^4$ $c = 16c - 48$ $c = 48/15 = 16/5$	M1	Equate difference in y -coordinates to ± 4
			M1	Use $\log a - \log b = \log a/b$
			A1	Obtain ${}^c/c-3 = 2^4$
			A1	Obtain $16/5$ oe
			[4]	

Q7, (OCR 4722, Jun 2013, Q8)

(i)	(a)	(0, 1)	B1	State (0, 1)
			[1]	
	(b)	(0, 4)	B1	State (0, 4)
			[1]	
	(c)	State a possible value for a	B1	Must satisfy $a > 1$
		State a possible value for b	B1	Must satisfy $0 < b < 1$
			[2]	

(ii)

$$\log_2 a^x = \log_2(4b^x)$$

$$\log_2 a^x = \log_2 4 + \log_2 b^x$$

$$x \log_2 a = \log_2 4 + x \log_2 b$$

$$x \log_2 a = \log_2 4 + x \log_2 (2/a)$$

$$x \log_2 a = 2 + x \log_2 2 - x \log_2 a$$

$$x (2 \log_2 a - 1) = 2$$

$$x = \frac{2}{2 \log_2 a - 1} \quad \mathbf{AG}$$

M1

Equate a^x and $4b^x$ and introduce logarithms at some stage

M1

Use $\log ab = \log a + \log b$ correctly

M1

Use $\log a^b = b \log a$ correctly at least once

B1

Use $b = 2/a$ to produce a correct equation in a and x only

A1

Obtain given relationship with no wrong working

[5]

(a) <u>Either</u> : State proportion $\frac{440}{275}$	B1
Attempt calculation involving proportion	M1 [involving multn and X value]
Obtain 704	A1 3
<u>Or</u> : Use formula of form $275e^{kt}$ or $275a^t$	M1 [or equiv]
Obtain $k = 0.047$ or $a = \sqrt[10]{1.6}$	A1 [or equiv]
Obtain 704	A1 (3) [allow ± 0.5]
(b)(i) Attempt correct process involving logarithm	M1 [or equiv including systematic trial and improvement attempt]
Obtain $\ln \frac{20}{80} = -0.02t$	A1 [or equiv]
Obtain 69	A1 3 [or greater accuracy; scheme for T&I: M1A2]
(ii) Differentiate to obtain $ke^{-0.02t}$	M1 [any constant k different from 80]
Obtain $-1.6e^{-0.02t}$ (or $1.6e^{-0.02t}$)	A1 [or unsimplified equiv]
Obtain 0.88	A1 3 [or greater accuracy; allow -0.88]

Q9, (OCR 4723, Jun 2008, Q7)

(i) State $A = 42$	B1
State $k = \frac{1}{9}$	B1 or 0.11 or greater accuracy
Attempt correct process for finding m	M1 involving logarithms or equiv
Obtain $\frac{1}{9} \ln 2$ or 0.077	A1 or 0.08 or greater accuracy
	4
(ii) Attempt solution for t using either formula	M1 using correct process (log'ms or T&I or ...)
Obtain 11.3	A1 or greater accuracy; allow 11.3 ± 0.1
	2
(iii) Differentiate to obtain form Be^{mt}	M1 where B is different from A
Obtain $3.235e^{0.077t}$	A1 ✓ or equiv; following their A and m
Obtain 47.9	A1 allow 48 or greater accuracy
	3

Q10, (OCR 4723, Jan 2009, Q5)

(i)	State 40	B1	
	Attempt value of k using 21 and 80	M1	or equiv
	Obtain $40e^{21k} = 80$ and hence 0.033	A1	or equiv such as $\frac{1}{21} \ln 2$
	Attempt value of M for $t = 63$	M1	using established formula or using exponential property
	Obtain 320	A1	5 or value rounding to this

(ii)	Differentiate to obtain $ce^{0.033t}$ or $40ke^{kt}$	M1	any constant c different from 40
	Obtain $40 \times 0.033e^{0.033t}$	A1√	following their value of k
	Obtain 2.64	A1	3 allow 2.6 or 2.64 ± 0.01 or greater accuracy (2.64056...)

8

Q11, (OCR 4723, Jan 2012, Q7)

(i)	(a)	State or imply $e^{-0.132t} = 0.25$ Attempt solution of eqn of form $e^{-0.132t} = k$ Obtain 10.5	B1 M1 A1 [3]	or equiv such as $40e^{-0.132t} = 10$ using sound process; implied by correct ans; allow trial and improvement attempt or greater accuracy
(i)	(b)	Differentiate to obtain $ke^{-0.132t}$ Obtain $5.28e^{-0.132t}$ or $-5.28e^{-0.132t}$ Substitute 5 to obtain 2.73 or -2.73	M1 A1 A1 [3]	where k is a constant not equal to 40 (allow even if process looks like integration) or (unsimplified) equiv accept 2.7 or -2.7 or greater accuracy; allow 2.73 or -2.73 whatever it is claimed to be
(ii)		<u>EITHER</u> Attempt to solve $40e^{2\lambda} = 31.4$ or $40e^{-2\lambda} = 31.4$ Obtain or imply $40e^{-0.121t}$ Substitute 3 to obtain 27.8 <u>OR</u> Attempt calculation involving multiplication of power of $\frac{31.4}{40}$ Obtain $31.4 \times (\frac{31.4}{40})^{0.5}$ or $40 \times (\frac{31.4}{40})^{1.5}$ Obtain 27.8	M1 A1 A1 [3] M1 A1 A1	using sound process; method implied by correct formula for mass of B obtained or greater accuracy ($-0.12103..$) or $0.5 \ln 0.785$ accept 28 or greater accuracy accept 28 or greater accuracy

Q12, (OCR 4723, Jun 2014, Q5)

(a)	Differentiate to produce $ke^{-0.33t}$ Obtain $-19.14e^{-0.33t}$ or $19.14e^{-0.33t}$ Obtain -5.1 or 5.1	M1 A1 A1 [3]	where constant k is different from 58 or unsimplified equiv whatever they claim value represents; accept 5.11 but not greater accuracy
(b)	<u>Either:</u> State or imply formula $42e^{kt}$ or $42a^t$ Attempt to find k from $42e^{6k} = 51.8$ or a from $42a^6 = 51.8$ Obtain $k = 0.035$ or $a = 1.0356$ Substitute 24 to obtain value between 97.1 and 97.3 inclusive	B1 M1 A1 A1	$42e^{-kt}$, $42e^{-kx}$, etc. also acceptable using sound process involving logarithms at least as far as $6k = \dots$ or $a = \dots$ or greater accuracy 0.03495... or exact equiv $\frac{1}{6} \ln \frac{37}{30}$ allow greater accuracy than 3 s.f.
	<u>Or:</u> Use ratio $\frac{51.8}{42}$ in calculation Attempt calculation of form $42 \times r^n$ Obtain $42 \times (\frac{51.8}{42})^4$ or $51.8 \times (\frac{51.8}{42})^3$ Obtain value between 97.1 and 97.3 inclusive	B1 M1 A1 A1 [4]	allow greater accuracy than 3 s.f.

Q13, (OCR 4723, Jun 2016, Q3)

i	Obtain 128 for value corresponding to 10 Obtain 65.5 for value corresponding to 25	B1 B1 [2]	Allow any value rounding to 128 Allow any value rounding to 65 or 66; whether obtained using powers of 0.8 or by use of formula
ii	Attempt to find formula for m of form $200e^{kt}$ or $200 \times r^{\lambda t}$ Obtain $200e^{(0.2 \ln 0.8)t}$ or $200e^{-0.0446t}$ or $200 \times 0.8^{0.2t}$ or 200×0.956^t Show correct process for solving equation of form $200e^{kt} = 50$ or $200r^{\lambda t} = 50$ Obtain 31	M1 A1 M1 A1 [4]	Whether attempted in part (i) or (ii) Or equiv Or greater accuracy rounding to 31; ignore any units given; second M1 is implied by correct answer